

This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package label for complete use directions. This product may not yet be available or approved for sale or use in your area.

# ZoGuard<sup>®</sup> plus FOR DOGS

## READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.

ZoGuard<sup>™</sup> Plus for Dogs contains Fipronil and S-Methoprene (insect growth regulator) to effectively kill fleas and ticks (including those that may transmit Lyme disease). The active ingredients in ZoGuard Plus for Dogs feature a sustained release formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for up to 1 month on dogs.

### ACTIVE INGREDIENTS:

Fipronil: .....	9.80%
(S)-methoprene: .....	8.80%
<b>OTHER INGREDIENTS:</b> .....	81.40%
<b>TOTAL:</b> .....	100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS. CAUTION.** Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**HAZARDS TO DOMESTIC ANIMALS.** For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

### FIRST AID

<b>If swallowed:</b>	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
<b>If in eyes:</b>	Flush eyes with plenty of water. Call a doctor if irritation persists.
<b>If on skin:</b>	Wash with plenty of soap and water. Get medical attention if irritation persists.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.** Do not use on cats, rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 8 weeks of age. Use entire contents of tube vial on each dog. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

**How to open:** Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the lines.

**How to apply:** Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks.

Apply one tube as a spot on the dog's back between the shoulder blades.

**Side Effects:** Monitor your dog after application. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-844-685-9173. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

### FREQUENCY OF APPLICATION

Use ZoGuard Plus for Dogs monthly for complete control of flea, tick, and chewing lice infestations. Studies show that ZoGuard Plus for Dogs kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. ZoGuard Plus for Dogs remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

If you have questions or comments about this product, please write:  
Promika, LLC

501 Cascade Pointe Lane, Suite 102, Cary, NC 27513

©2014 Promika. ZoGuard<sup>®</sup> is a trademark of Promika, LLC.

Frontline<sup>®</sup> is a registered trademark of Merial. ZoGuard<sup>®</sup> Plus for Dogs is not manufactured, or distributed by Merial, seller of Frontline<sup>®</sup> Plus for Dogs.



**DO NOT USE ON CATS.** Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.

Contains Fipronil and S-Methoprene,  
the active ingredients used in  
FRONTLINE<sup>®</sup> Plus for Dogs

**Only For Dogs Weighing 4 to 22 lbs.**  
Do not apply to dogs weighing less than 4 lbs.  
including small dogs and puppies, 8 weeks or older:  
Apply one tube (0.67 mL) (0.023 fl oz)

Three 0.023 fl oz (0.67 mL) Applicators  
Net Contents: 0.069 fl oz

**Only For Dogs Weighing 23-44 lbs.**  
Apply one tube (1.34 mL) (0.045 fl oz)






Three 0.045 fl oz (1.34 mL) Applicators  
Net Contents: 0.135 fl oz

**Only For Dogs Weighing 45 to 88 lbs.**  
Apply one tube (2.68 mL) (0.091 fl oz)

Three 0.091 fl oz (2.68 mL) Applicators  
Net Contents: 0.273 fl oz

**Only For Dogs Weighing 89 to 132 lbs.**  
Apply one tube (4.02 mL) (0.136 fl oz)

Three 0.136 fl oz (4.02 mL) Applicators  
Net Contents: 0.408 fl oz

- KILLS**  **FLEAS**
-  **TICKS**
-  **FLEA EGGS**
-  **FLEA LARVAE**
-  **CHEWING LICE**

### DIRECTIONS FOR USE

#### How to Open

1. Remove product tubes from the package.
2. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.

#### How to Apply

1. Invert tube over dog and use open end to part dog's hair.
2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot on the dog's back between the shoulder blades.



### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight

**CONTAINER DISPOSAL:** If empty: Non-refillable container. Do not reuse or refill this container.

Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Distributed by:  
**Promika, LLC.**  
501 Cascade Pointe Lane, Suite 102  
Cary, NC 27513

In Case of Emergency,  
Call: 1-844-685-9173

EPA Reg. No. 53883-351-91300