

This information is for promotional purposes only. Space considerations may require information to be omitted. Always refer to the actual package label for complete use directions. This product may not yet be available or approved for sale or use in your area.

Promika ZoGuard[®] plus FOR DOGS

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND PUPPIES 5 LBS AND OVER & 8 WEEKS OF AGE OR OLDER.**

Promika ZoGuard[®] Plus for Dogs contains Fipronil and (S)-Methoprene (insect growth regulator) to effectively kill fleas and ticks including those that may transmit Lyme disease. The active ingredients in Promika ZoGuard Plus for Dogs features a formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for up to 1 month on dogs.

ACTIVE INGREDIENTS:

Fipronil:.....	9.80%
(S)-methoprene:	8.80%
OTHER INGREDIENTS:	81.40%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product. This product is not a substitute for prescribed Heartworm preventative.

FIRST AID	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information, you may contact SafetyCall at 1-844-685-9173 seven days a week, 24-hours a day.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs.

How to open: Refer to package insert for directions on how to open package.

How to apply: Remove product applicator from the package. Separate one applicator from the others. Hold the applicator with notched end pointing up and away from the face and body. Snap the outline of the tip back to allow access to the dotted tip of applicator. Holding the applicator away from your and your dog's eyes, cut applicator tip with scissors on dotted line. Invert applicator over dog and use open end to part dog's hair. Squeeze applicator firmly to apply all of the solution to the dog's skin as a spot on to the dog's back between the shoulder blades. Allow area to dry. Avoid contact with treated area until dry. Use only one applicator per treatment.

Side Effects (Dogs): Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. In some cases gastrointestinal signs such as vomiting and diarrhea may occur. If these side effects occur, consult your veterinarian or call SafetyCall at 1-844-685-9173. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application.

PRODUCT INFORMATION

Use Promika ZoGuard Plus for Dogs monthly for complete control of flea, tick, and chewing lice infestations. Studies show that this product kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. Promika ZoGuard Plus for Dogs remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Avoid contact with treated area until dry. Do not reapply for 30 days.

If you have questions or comments about this product, please write:

Promika, LLC
5000 CentreGreen Way, Suite 100, Cary, NC 27513

©2018 Promika. ZoGuard[®] is a trademark of Promika, LLC. Frontline[®] is a registered trademark of Merial. ZoGuard[®] Plus for Dogs is not manufactured, or distributed by Merial, seller of Frontline[®] Plus for Dogs.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.


**Contains Fipronil and (S)-Methoprene,
the active ingredients used in
FRONTLINE[®] Plus for Dogs**

**Only For Dogs Weighing 5 to 22 lbs.
& 8 weeks or older. Do not apply to dogs
or puppies weighing less than 5 lbs.**
Apply one applicator (0.67 mL) (0.023 fl oz)

**Only for Dogs Weighing 23 to 44 lbs.
& 8 weeks and older.**
Apply one applicator (1.34 mL) (0.045 fl oz)

**Only for Dogs Weighing 45 to 88 lbs.
& 8 weeks and older.**
Apply one applicator (2.68 mL) (0.091 fl oz)

**Only for Dogs Weighing 89 to 132 lbs.
& 8 weeks and older. Do not use more than one
applicator per treatment for one dog regardless
of the dog's weight.**
Apply one applicator (4.02 mL) (0.136 fl oz)

- KILLS**  **FLEAS**
-  **TICKS**
-  **FLEA EGGS**
-  **FLEA LARVAE**
-  **CHEWING LICE**

DIRECTIONS FOR USE

Remove product applicator from the package. Separate one applicator from the others. Hold the applicator with notched end pointing up and away from the face and body. Snap the outline of the tip back to allow access to the dotted tip of applicator. Holding the applicator away from your and your dog's eyes, cut applicator tip with scissors on dotted line. Invert applicator over dog and use open end to part dog's hair. Squeeze applicator firmly to apply all of the solution to the dog's skin as a spot on to the dog's back between the shoulder blades. Allow area to dry. Avoid contact with treated area until dry. Use only one applicator per treatment.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not remove applicator from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight

PESTICIDE DISPOSAL AND CONTAINER HANDLING:
If empty: Offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Distributed by:
Promika, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

In Case of Emergency,
Call: 1-844-685-9173

EPA Reg. No. 91300-5